Patient Name:

PROHIBITED MEDICATIONS/SUBSTANCES TO STOP BEFORE IN-OFFICE PROCEDURE

Please STOP the following medications before and after your surgery as specified in the table below. These medications have been known to interfere with surgery, the amount of bleeding and bruising during or after your surgery and delay the healing process. Please keep in mind this IS NOT a comprehensive list so if you have ANY QUESTIONS AS TO A MEDICATION YOU ARE CURRENTLY TAKING, CALL US IMMEDIATELY FOR GUIDANCE. If there is a medication listed that your doctor does NOT want you to stop, please notify us immediately. TYLENOL IS FINE TO TAKE.

Please alert us if you are taking Semiglutides as your instructions for before the procedure may change.

Stop 2 weeks Before & After	Stop 30 Days Before & After	Stop 6 Weeks Before & After
<u>Procedure</u>	<u>Procedure</u>	<u>Procedure</u>
Alcohol	Betamethasone	Green Tea/Herbal Teas
Aspirin	Depo-Medrol	Multivitamins
Aspirin Containing Products	Marijuana	
Advil	Medrol Dosepack	
Aleve	Methyl Prednisolone	
Alka-Seltzer products	Nicotine	
Aspergum	Prednisone	
Celebrex	Prednisolone	
Chia seeds	Smoking	
Diclofenac	Solumedrol	
Excedrin	Steroids/Glucosteroids	
Fish Oil	Vaping	
Flax Seed		
Ibuprofen		
Indomethacin		
Meloxicam		
Mobic		
Motrin		
Naproxen		
Omegas		
Percodan		
Phenedimetrazine		
Phentermine		
St. John's Wort		
Valeriean		
Vitamin E		
Voltaren		
		9199
-		

	SIS!			
I have received, read, and understand this document.				
Patient Signature	Date			