Patient Name:

PROHIBITED MEDICATIONS/SUBSTANCES TO STOP BEFORE SURGERY

<u>Please STOP the following medications before and after your surgery as specified in the table below.</u> These medications have been known to interfere with surgery, the amount of bleeding and bruising during or after your surgery and delay the healing process. Please keep in **mind this IS NOT a comprehensive list so if you have ANY QUESTIONS AS TO A MEDICATION YOU ARE CURRENTLY TAKING, CALL US IMMEDIATELY FOR GUIDANCE.** If there is a medication listed that your doctor does NOT want you to stop, please notify us immediately. TYLENOL IS FINE TO TAKE.

STOP 24-48 HOURS BEFORE SURGERY	Stop 2 weeks Before & After Surgery	Stop 30 Days Before & After Surgery	Stop 6 Weeks Before & After Surgery
Any blood pressure	Alcohol	Betamethasone	Green Tea/Herbal Teas
medication			
Metoprolol	Aspirin	Depo-Medrol	Mounjaro
Valsartan HCTZ	Aspirin Containing Products	Marijuana	Multivitamins
Lisinopril	Advil	Medrol Dosepack	Ozempic
Amlodipine	Aleve	Methyl Prednisolone	Semiglutides
	Alka-Seltzer products	Nicotine	Wegovy
	Aspergum	Prednisone	Weight Loss Medications
	Celebrex	Prednisolone	Weight Loss Injections
	Chia seeds	Smoking	
	Diclofenac	Solumedrol	
	Excedrin	Steroids/Glucosteroids	
	Fish Oil	Vaping	
	Flax Seed	-	
	Ibuprofen		
	Indomethacin		
	Meloxicam		
	Mobic		
	Motrin		
	Naproxen		
	Omegas		
	Percodan		
	Phenedimetrazine		
	Phentermine		
	St. John's Wort		
	Valeriean		
	Vitamin E		
	Voltaren		

I have received, read, and understand this document.				
Patient Signature	Date			